



PICKLING, SLICING & SPECIALTY CUCUMBER VARIETIES COMPARISON CHART



Dimensions may vary significantly due to culture and environmental conditions.

Part Number	Variety	Days to Maturity	Harvest Length	Disease Resistance	Gynoecious	Parthenocarpic
Pickling						
2563G OG	Adam Gherkin	47	2–3"	(IR: CMV, PM, S.)	x	x
3541	Citadel	52	4–5"	(HR: A, ALS, S. IR: CMV, DM, PM.)	x	
3449G OG	Cool Customer	55	4–5"			
3082G OG	Excelsior	50	4–5"	(HR: S, TSP. IR: CMV, CVYV, PM.)	x	x
331G OG	H-19 Little Leaf	57	3–5"	(HR: A, ALS, B, S. IR: CMV, PM.)		x
329G OG	Lemon	65	1½–2½"			
3346	Max Pack	55	4–5"	(HR: A, S, WMV, ZYMV. IR: ALS, CMV, PM, PRSV.)		
330	Northern Pickling	48	3–5"	(HR: S.)		
2524G OG	Salt and Pepper	49	3–5"	(IR: ALS, PM.)		
Seedless/Thin Skinned						
2198 or 2198T	Diva	58	5–7"	(HR: S. IR: CVYV, PM.)		x
3114	Iznic	45	3½–4"	(HR: PM, S.)	x	x
Slicing						
3540	Bristol	54	8"	(HR: A, ALS, CMV, S, ZYMV. IR: DM, PM, PRSV.)	x	
2562G OG	Corinto	48	7–8"	(IR: CMV, CVYV, PM.)	x	x
4536T or 2417	General Lee	52	8–8½"	(HR: S. IR: CMV, PM.)	x	
336G OG , or 336	Marketmore 76	58	8–9"	(HR: S. IR: CMV, PM.)		
2903T or 2903	Olympian	52	8–9"	(IR: A, ALS, PM, S.)	X	
2323	Tasty Jade	54	11–12"	(IR: PM.)		X
Specialty						
3436	Itachi	54	9–11"			x
3174	Mexican Sour Gherkin	67	1"			
3524	Nokya	55	11–12"	(IR: PM.)		x
2317	Striped Armenian	63	8–18"			
333G OG	Suyo Long	61	10–15"			

Part Numbers with T=Treated seeds. **OG** = Organic seeds.

Cucumber Resistance Codes

HR: = High Resistance IR: = Intermediate Resistance

(A)	Anthraxnose
(ALS)	Angular Leaf Spot
(B)	Bacterial Wilt
(CMV)	Cucumber Mosaic Virus
(CVYV)	Cucumber Vein Yellowing Virus
(DM)	Downy Mildew
(PM)	Powdery Mildew
(PRSV)	Papaya Ringspot Virus
(S)	Scab
(TSP)	Target Spot
(WMV)	Watermelon Mosaic Virus
(ZYMV)	Zucchini Yellow Mosaic Virus

GROWING DEFINITION

Gynoecious: Produces only female flowers but must be pollinated by males to set fruit. For this reason, gynoecious varieties are blended with 10–15% seed of monoecious cucumbers. Produces in a more concentrated period. Preferred for mechanical harvest, or other situations where less-frequent harvesting is desired.

Parthenocarpic: Does not require pollination to set fruit. When successfully isolated from pollen, their fruits will be seedless. Can be either monoecious or gynoecious and are recommended for use in protected culture.